

ELLIPTIC UNITS IN K_2

KENICHIRO KIMURA

Institute of Mathematics, University of Tsukuba, Tsukuba,
Ibaraki, 305 Japan email: kimurak@math.tsukuba.ac.jp

abstract: We present certain norm-compatible systems in K_2 of function fields of some CM elliptic curves. We demonstrate that these systems have some properties similar to elliptic units.

Key words: CM elliptic curve, K_2 , Coleman power series, explicit reciprocity law, Euler system

§1 Introduction. Elliptic units are systems of units in abelian extensions of imaginary quadratic fields which are defined by evaluating some functions on CM elliptic curves at torsion points. These units play an essential role in Iwasawa theory of imaginary quadratic fields ([dS],[Ru] etc). In this paper we present certain systems of K_2 of function fields of CM elliptic curves. These systems are considered by Scholl ([Sch]) in a more general setting. We show that these systems have two similar properties to elliptic units.

First, in §2 we demonstrate that we can apply the theory of Coleman power series for K_2 of 2-dimensional local fields constructed by Fukaya ([Fu]) to our system. We show that the “Coleman power series” of our systems, which are elements in K_2 of some power series rings, are explicitly described. In §3 we modify our systems so that they are in the kernel of tame symbol maps. Using the result of §2 we will give a sufficient condition for the images of those systems to be non-zero under Chern class maps to Galois cohomology groups.

Second, in the appendix we show that our systems form Euler systems similar to those of elliptic units.

By [Sch] one can see that systems like ours also exist in higher K -groups of function fields of products of elliptic curves. This fact seems to suggest (to this author) that there is a further generalization of the theory of elliptic units and Coleman power series to higher K -groups.

¹The author was partially supported by JSPS Postdoctoral Fellowships for Research Abroad from 2000-2001.

The following question is asked by the referee:

Is there interpretation of the function which appears in corollary 3.4 in terms of L -function?

Acknowledgment. The author would like to express his gratitude to Prof. K. Kato for valuable suggestions and to Dr.T. Fukaya for showing her paper at an early stage. He is also grateful to the referee for valuable comments on the first version of this paper. This work was carried out while the author stayed at Department of Mathematics, the University of Chicago. He is thankful to the people at Chicago for their hospitality.

Notations.

K :	an imaginary quadratic field,	4
E :	a CM elliptic curve,	4
F :	an abelian extension of K ,	4
ψ :	the Hecke character of F associated to E ,	5
ϕ :	a Hecke character of K such that $\psi = \phi \circ N_{F/K}$,	5
$\mathfrak{f}$:	conductor of E ,	5
p :	a rational prime,	5
$\mathfrak{p}$:	a prime factor of p in $\mathcal{O}_K$,	5
π :	$\phi(\mathfrak{p})$,	5
a :	an integer relatively prime to $6p\mathfrak{f}$,	5
g_a :	a rational function on E ,	5
v_n :	a generator of $E[\pi^n]$,	5
T_y :	translation by a point y ,	5
F_n :	$F(E[\pi^n])$,	5
α_n :	an element of $K_2(F_n(E))$,	5
W :	the Witt ring of $\overline{\mathbb{F}_p}$,	6
$\mathfrak{P}$:	a prime of F over p ,	6
$\hat{\mathcal{E}}_W$:	the formal completion of a projective smooth model of E ,	6
θ :	an isomorphism $\widehat{\mathbb{G}_{mW}} \rightarrow \hat{\mathcal{E}}_W$,	6
Φ :	Frobenius of W ,	6
t :	a parameter on $\hat{\mathcal{E}}_W$,	6
s :	the standard parameter on $\widehat{\mathbb{G}_{mW}}$,	6
$(\zeta_n)_{n \leq 1}$:	a basis of $\mathbb{Z}_p(1)$,	6
H :	p -adic completion of $W((s))$,	6
H_n :	$H(s_n, \zeta_n)$,	6
S :	$\mathcal{O}_H[[\epsilon - 1]]$,	6
S' :	$S[(\epsilon - 1)^{-1}]$,	6

- ν, σ, φ : endomorphisms of S , 6
- h : an isomorphism $K_2(S')^{\wedge^{Nr_\varphi=1}} \simeq \varprojlim_n K_2(H_n)^\wedge$, 7
- G_k : $\text{Gal}(k^{sep}/k)$ for a field k , 7
- $H^i(k, M)$: $H^i(G_k, M)$ for a G_k module M , 7
- λ, ∂ : homomorphisms $\varprojlim_n K_2(H_n)^\wedge \rightarrow \Omega_{H_1}^1$, 7
- $q : s + 1$, 8
- β_n : $\theta^{\Phi^{-n}*} \alpha_n$, 8
- D : $q \frac{d}{ds}$, 9
- F'_n : $F(E[a\pi^n])$, 9
- α_n^γ : $T_\gamma^* \alpha_n$, 9
- α'_n : an element of $K_2(F_n(E))$, 9
- β'_n : $\theta^{\Phi^{-n+1}*} \alpha'_n$, 10
- g'_a : $T_\gamma^* g_a$, 11

§2 norm-compatible systems in K_2 .

A general reference for this section is Chapter II of [dS]. We fix an imaginary quadratic field K . Let O_K denote the ring of integers of K . Let F be an abelian extension of K and let E be an elliptic curve over F which satisfies the following conditions:

- (i) E has complex multiplication by O_K , and
- (ii) $F(E_{tors})$ is an abelian extension of K .

Here $F(E_{tors})$ denotes the field generated by coordinates of torsion points of a model of E over F . If ψ is the Hecke character of F associated to E , then the condition (ii) is equivalent to the existence of a Hecke character ϕ of K such that $\psi = \phi \circ N_{F/K}$.

Let $\mathfrak{f}$ be the conductor of E and let $p > 3$ be a rational prime which is relatively prime to $\mathfrak{f}$ and which splits completely in F . Let $p = \mathfrak{p}\bar{\mathfrak{p}}$ be the decomposition of p in O_K . We write $\pi = \phi(\mathfrak{p})$. Note that $\mathfrak{p} = (\pi)$. We assume that $\text{Gal}(F(E[p^n]))/F \simeq (O_K/(p^n))^*$ for all $n \geq 1$.

Let a be an integer such that $(a, 6p\mathfrak{f}) = 1$. Then there is a function $g_a \in F(E)$, which is called the Kato-Siegel function in [Sch]. We now state the properties of g_a which will be used in the subsequent discussions.

- (a) $\text{div} g_a = a^2(0) - E[a]$.
- (b) For any isogeny $j : E \rightarrow E$ defined over $\bar{F}$ such that $(\deg j, a) = 1$, $j_* g_a = g_a$.
- (c) $[a]_* g_a = 1$.
- (d) $[-1]^* g_a = g_a$. (This is a special case of (b)).

Here for an element u of O_K , $[u]$ stands for an element of $\text{End}(E)$, which is justified by the condition (i). In our situation, the property (b) is equivalent to the following

distribution relation

$$\prod_{j(y)=x} g_a(y) = g_a(x).$$

We fix a system $(v_n)_{n \geq 1}$ of torsion points of $E(\bar{F})$ such that for each $n \geq 2$, v_n is a generator of $E[\pi^n]$ and that $[\pi]v_n = v_{n-1}$.

For a point y on E we denote by $T_y : E \rightarrow E$ the translation by y . For $n \geq 1$ consider the element

$$\alpha_n := \{g_a, T_{v_n}^* g_a\} \in K_2(F_n(E)).$$

Here $F_n := F(E[\pi^n])$ and $F_n(E)$ is the function field of $E_{F_n} = E \times_F F_n$. This element is considered by Scholl([Sch], Appendix) in a more general setting. The following lemma claims that the system (α_n) forms an norm-compatible system.

Lemma 2.1. *For $n \geq 2$, the equality*

$$[\pi]_* N_{F_n(E)/F_{n-1}(E)} \alpha_n = \alpha_{n-1}$$

holds.

Proof. This is another way of describing (EN1) in loc.cit. We will give a proof by calculation of symbols.

$$N_{F_n(E)/F_{n-1}(E)} \alpha_n = \{g_a, \prod_{v \in E[\pi]} T_{v_n+v}^* g_a\} = \{g_a, [\pi]^*(T_{v_{n-1}}^* g_a)\}$$

where the second equality holds by the distribution relation mentioned above, and

$$[\pi]_* \{g_a, [\pi]^*(T_{v_{n-1}}^* g_a)\} = \{[\pi]_* g_a, T_{v_{n-1}}^* g_a\} = \{g_a, T_{v_{n-1}}^* g_a\}.$$

We will apply the theory of Coleman power series for K_2 constructed by Fukaya [Fu] to this system. As at present her theory treats only the multiplicative group, we need to construct an isomorphism from E to the multiplicative group. For this we will take the formal completion of E along the 0-section.

We will follow the argument of §4.3, Chapter II of [dS] rather closely. Let $F' = F(E[\pi^\infty])$ and let R be the completion of $O_{F'} \otimes_{O_K} O_{K_{\mathfrak{p}}}$. Since the prime $\mathfrak{p}$ finitely decomposes in F' , R is a direct sum of local rings. Fix a direct summand of R and denote it by W . Let $\mathfrak{P}$ be the prime of F over $\mathfrak{p}$ which corresponds to W and let $\mathcal{E}$ be a smooth projective model of E over $O_{F_{\mathfrak{P}}} = \mathbb{Z}_p$. Let $\hat{\mathcal{E}}_W$ be the formal completion of $\mathcal{E} \times_{\mathbb{Z}_p} W$ along the 0-section. Then by Proposition 4.3 in Chap.2 of [dS], there is an isomorphism

$$\theta : \widehat{\mathbb{G}_{mW}} \rightarrow \hat{\mathcal{E}}_W$$

which satisfies the equality $[\pi] \circ \theta = \theta^\Phi \circ [p]$. Here Φ is the Frobenius automorphism of W . We fix a parameter t of $\hat{\mathcal{E}}_W$ and let s be the standard parameter of $\widehat{\mathbb{G}_{mW}}$. For $n \geq 1$, define $\zeta_n \in \overline{\text{Frac}(W)}$ by the equality $t(v_n) = \theta^{\Phi^{-n}}(\zeta_n - 1)$. Then we see that the system $(\zeta_n)_{n \geq 1}$ is a basis of $\mathbb{Z}_p(1)$.

Now we will review the theory of Fukaya in the form we will use in the subsequent discussions.

Let H be the p -adic completion of $W((s))$, the function field of $\widehat{\mathbb{G}_{mW}}$. Consider a tower of field extensions $(H_n)_{n \geq 1}$ where $H_n := H(s_n, \zeta_n)$ and $(s_{n-1} + 1) = (s_n + 1)^p$ for $n \geq 1$. $s_0 = s$ by convention.

We set $S := O_H[[\epsilon - 1]]$ to be the ring of formal power series and write $S' = S[(\epsilon - 1)^{-1}]$. Consider the ring homomorphisms ν , σ and $\varphi : S' \rightarrow S'$ defined as follows:

ν is defined by $\nu(\epsilon) = \epsilon^p$ and $\nu|_{O_H} = \text{the identity}$. The homomorphism σ is defined by $\sigma|_{O_H} = [p]^* \otimes \Phi$ and $\sigma(\epsilon) = \epsilon$. The homomorphism φ is defined by the composition: $\varphi = \nu \circ \sigma$. The following theorem is a part of Theorem 1.5 of [Fu]. It asserts the existence of Coleman power series:

Theorem 2.2. *There is a canonical isomorphism*

$$h : K_2(S')^{\wedge N_{r_\varphi}=1} \simeq \varprojlim_n K_2(H_n)^\wedge$$

induced by

$$h_n : S' \rightarrow H_n, f(\epsilon) \mapsto (\sigma^{-n} f)(\zeta_n).$$

Here N_{r_φ} is the norm map on $K_2(S')$ induced by the finite morphism φ and $*^{N_{r_\varphi}=1}$ means the fixed part of $*$ under N_{r_φ} . The map $\sigma^{-n} : H \hookrightarrow H_n$ is the composition of $\text{id} \otimes \Phi^{-n} : H \rightarrow H$ and the inclusion given by $s \mapsto s_n$. The symbol $\wedge$ means certain completion.

Also there is a fact which corresponds to the explicit reciprocity law. It asserts the coincidence with $-$ sign of the two homomorphisms λ and ∂ defined as follows.

For a field k , we write G_k for $\text{Gal}(k^{sep}/k)$. For a G_k module M , we write $H^i(k, M)$ for $H^i(G_k, M)$.

$$\begin{aligned}
\lambda : \varprojlim_n K_2(H_n)^\wedge &\xrightarrow{cl} \varprojlim_n H^2(H_n, (\mathbb{Z}/p^n\mathbb{Z})(2)) \\
&\xrightarrow{\cup(\zeta_n)^{-1}} \varprojlim_n H^2(H_n, (\mathbb{Z}/p^n\mathbb{Z})(1)) \\
&\xrightarrow{Cor_{H_n/H_1}} \varprojlim_n H^2(H_1, (\mathbb{Z}/p^n\mathbb{Z})(1)) \\
&= H^2(H_1, \mathbb{Z}_p(1)) \\
&\rightarrow H^2(H_1, \mathbb{C}_p(1)) \simeq \Omega_{H_1}^1.
\end{aligned}$$

The last isomorphism is due to Hyodo [Hy]. Here $\Omega_{H_1}^1$ is defined to be $\mathbb{Q} \otimes \Omega_{O_{H_1}}$ where $\Omega_{O_L}^1 := \varprojlim_n \Omega_{O_{H_1}/\mathbb{Z}}^1 / p^n \Omega_{O_{H_1}/\mathbb{Z}}^1$.

The homomorphism ∂ is defined using the map h in Theorem 2.2:

$$\begin{aligned}
\partial : \varprojlim_n K_2(H_n)^\wedge &\xrightarrow{h^{-1}} K_2(S')^{Nr_\varphi=1} \\
&\xrightarrow{d\log} \Omega_S^2(\log) = \frac{S}{\epsilon - 1} d\log(q) \wedge d\log(\epsilon) \\
&\xrightarrow{(\wedge d\log(\epsilon))^{-1}} \frac{S}{\epsilon - 1} d\log(q) \\
&\rightarrow \Omega_{H_1}^1
\end{aligned}$$

where $q = s + 1$ and the last map is defined by

$$f(\epsilon) d\log(q) \mapsto \frac{1}{p} (\sigma^{-1} f)(\zeta_1) d\log(q).$$

The modules $\Omega_S^1(\log)$ and $\Omega_S^2(\log)$ are defined as follows:

Let

$$\Omega_{S/\mathbb{Z}}^1(\log) := (\Omega_{S/\mathbb{Z}}^1 \oplus S \otimes_{\mathbb{Z}} S'^*)/N$$

where N is the S submodule generated by elements $(-da, a \otimes a)$ for $a \in S \cap S'^*$.

The class $(0, 1 \otimes a)$ is denoted by $d\log(a)$. The module $\Omega_S^2(\log)$ is defined by

$$\Omega_S^2(\log) = \varprojlim_n \wedge^2 \Omega_{S/\mathbb{Z}}^1(\log) / p^n \wedge^2 \Omega_{S/\mathbb{Z}}^1(\log).$$

By proposition (19.3.5) of [Gr], S is formally smooth over $\mathbb{Z}$, so by corollary (20.4.11) of loc.cit. $\Omega_{S/\mathbb{Z}}^1(\log) / p^n \Omega_{S/\mathbb{Z}}^1(\log)$ is a free $S/p^n S$ module on $d\log(\epsilon - 1)$ and $d\log(q)$. Hence $\Omega_S^2(\log)$ is a free S module on $d\log(q) \wedge d\log(\epsilon - 1)$.

The map $d\log : K_2(S') \rightarrow \Omega_S^2(\log)$ is described as $\{f, g\} \mapsto d\log(f) \wedge d\log(g)$.

Theorem 2.3 (Theorem 2.4 in [Fu]).

$$\lambda = -\partial.$$

For $n \geq 1$, set $\beta_n := \theta^{\Phi^{-n}*} \alpha_n \in K_2(H_n)$. Since $[p]_* \theta^* = \theta^{\Phi^*} [\pi]_*$ it follows that $(\beta_n)_n$ is an element of $\varprojlim_n K_2(H_n)$ where projective limit is taken with respect to norm maps.

The following is our main result:

Theorem 2.4. *We have the equality*

$$h^{-1}((\beta_n)_n) = B := \{\theta^* g_a(s), \theta^* g_a(s[+]_{\widehat{\mathbb{G}_m W}} \epsilon - 1)\}.$$

Proof. First of all, note that since $g_a(t) \in t^{1-a^2} W[[t]]^*$, $\theta^* g_a(s) \in s^{1-a^2} W[[s]]^* \subset O_H^*$. Since $\theta^* g_a(s[+]_{\widehat{\mathbb{G}_m W}} \epsilon - 1)$ is of the form $\theta^* g_a(s) + (\epsilon - 1)f(s, \epsilon - 1)$ with $f(s, \epsilon - 1) \in S$, it belongs to S^* . It can also be checked directly that $h_n(B) = \beta_n$ for $n \geq 1$. What remains is to show that $Nr_\varphi B = B$.

$$\begin{aligned} Nr_\varphi B &= Nr_\sigma \circ Nr_\nu B \\ &= Nr_\sigma \circ \{\theta^* g_a, \prod_{\zeta^p=1} \theta^* g_a(s[+]_{\widehat{\mathbb{G}_m W}} \zeta \epsilon - 1)\} \\ &= Nr_\sigma \circ \{\theta^* g_a, \prod_{\zeta^p=1} \theta^* g_a(s[+]_{\widehat{\mathbb{G}_m W}} \epsilon - 1[+]_{\widehat{\mathbb{G}_m W}} \zeta - 1)\} \\ &= Nr_\sigma \circ \{\theta^* g_a, \prod_{\zeta^p=1} g_a(\theta(s[+]_{\widehat{\mathbb{G}_m W}} \epsilon - 1)[+]_{\widehat{\mathcal{E}_W}} \theta(\zeta - 1))\} \\ &= Nr_\sigma \circ \{\theta^* g_a, \prod_{v \in E[\pi]} g_a(\theta(s[+]_{\widehat{\mathbb{G}_m W}} \epsilon - 1)[+]_{\widehat{\mathcal{E}_W}} v)\} \\ &= Nr_\sigma \circ \{\theta^* g_a, g_a([\pi](\theta(s[+]_{\widehat{\mathbb{G}_m W}} \epsilon - 1)))\} \\ &= Nr_\sigma \circ \{\theta^* g_a, g_a(\theta^\Phi([p](s[+]_{\widehat{\mathbb{G}_m W}} \epsilon - 1)))\} \\ &= Nr_\sigma \circ \{\theta^* g_a, [p]^*(\theta^{\Phi^*} g_a)(s[+]_{\widehat{\mathbb{G}_m W}} \epsilon - 1)\}. \end{aligned}$$

Since $\sigma = [p]^* \otimes \Phi$, $Nr_\sigma = \Phi^{-1} \circ [p]_*$. From this and the equality $[p]_*(\theta^* g_a) = \theta^{\Phi^*} [\pi]_* g_a = \theta^{\Phi^*} g_a$ it follows that $Nr_\varphi B = B$. $\square$

Let D be a differential operator on H defined by $D = q \frac{d}{ds}$. Then by straightforward calculation we obtain the following corollary.

Corollary 2.5.

$$\partial((\beta_n)_n) = \frac{1}{p} \Phi^{-1}(D \log(\theta^* g_a)(s) \times D \log(\theta^* g_a)(s[+])_{\widehat{\mathbb{G}_{m_W}}} \zeta_1 - 1) d \log q.$$

§3. Elements in the kernel of tame symbol map.

For $n \geq 1$, let $F'_n := F(E[a\pi^n])$. For a point $\gamma \in E[a]$, let $\alpha_n^\gamma := T_\gamma^* \alpha_n \in K_2(F'_n(E))$. We set

$$\alpha'_n := N_{F'_n/F_n}[\pi]_* \sum_{\gamma \in E[a]} \alpha_n^\gamma.$$

We can check that for $n \geq 2$,

$$[\pi]_* N_{F_n(E)/F_{n-1}(E)} \alpha'_n = \alpha'_{n-1}$$

in a similar way as in the case of α_n . We have the following exact sequence ([Qu]):

$$\bigoplus_{x \in E(\bar{F})} K_2(\kappa_x) \rightarrow K_2(E_{F_n}) \xrightarrow{r} K_2(F_n(E)) \xrightarrow{\bigoplus_{x \in E(\bar{F})} t_x} \bigoplus_{x \in E(\bar{F})} \kappa_x^* \xrightarrow{N} F_n^*$$

where r is the restriction, N is the norm and for $x \in E(\bar{F})$, t_x is defined by

$$\{f, g\} \mapsto (-1)^{\text{ord}_x f \text{ord}_x g} \frac{f^{\text{ord}_x g}}{g^{\text{ord}_x f}}(x).$$

By this exact sequence we see that $[\pi^n]^* [\pi^n]_* \sum_{\gamma \in E[a]} \alpha_n^\gamma = \sum_{u \in E[a\pi^n]} T_u^* \alpha_n$ is in the kernel of $\bigoplus_{x \in E(\bar{F})} t_x$, and since the map $[\pi^n] : E \rightarrow E$ is etale, it follows that

$$\bigoplus_{x \in E(\bar{F})} t_x([\pi^{n-1}]_* \alpha'_n) = 0.$$

Hence there is an element $\mathcal{A} \in K_2(E_{F_n})$ such that $r(\mathcal{A}) = [\pi^{n-1}]_* \alpha'_n$. Since K_2 of number fields is torsion ([Ga]), $\mathcal{A}$ is well defined up torsion.

Lemma 3.1. *Let $V_p(E)(1) := T_p(E)(1) \otimes \mathbb{Q}$ and set $F_\infty := F(E[\pi^\infty])$. Then*

$$(V_p(E)(1)/T_p(E)(1))^{G_{F_\infty}} = 0.$$

Proof. There is a decomposition $T_p(E)(1) = T_{\mathfrak{p}}(E)(1) \oplus T_{\bar{\mathfrak{p}}}(E)(1)$, so we will treat each factor separately. The factor $T_{\mathfrak{p}}(E)(1)$ is isomorphic to $\mathbb{Z}_p(1)$ as a G_{F_∞} module. By corollary 1.7 in Chap.II of [dS], the fields F_∞ and $F(E[\bar{\pi}^\infty])$ are linearly disjoint

over F . Since $\mathbb{Z}_p(1) \simeq T_{\mathfrak{p}}(E) \otimes T_{\bar{\mathfrak{p}}}(E)$, the fields F_{∞} and the field $F(\zeta_{p^{\infty}})$, generated by p -power roots of unity, are also linearly disjoint over F . Hence the assertion holds for this factor.

As for $T_{\bar{\mathfrak{p}}}(E)(1)$, it is isomorphic to $\mathbb{Z}_p(2)$ as a $G_{F_{\infty}}$ module. Since $p > 3$, the assertion holds for this factor as well. $\square$

By the lemma, $H^1(F_n, T_p(E)(1))$ has no torsion and we have a well defined class $\tilde{cl}([\pi^{n-1}]_* \alpha'_n) := cl(\mathcal{A}) \in H^1(F_n, T_p(E)(1))$.

We set $\beta'_n := \theta^{\Phi^{-n+1}*} \alpha'_n \in K_2(H_n)$. Since all the points in $E[a]$ are rational over $\text{Frac} W$ and $\theta^{\Phi*}[\pi]_* = [p]_* \theta^*$, we have the equality

$$\beta'_n = [F'_0 : F][p]_* \sum_{\gamma \in E[a]} \theta^{\Phi^{-n}*} \alpha_n^{\gamma}.$$

One can check that $(\beta'_n)_n$ form an element of $\varprojlim_n K_2(H_n)$ in a similar way to the case of β_n .

It can also be shown that $\theta^*([\pi^{n-1}]_* \alpha'_n) = [p^{n-1}]_* \beta'_n$. By (5.10) in [Fu], we have a commutative diagram

$$\begin{array}{ccccc} \varprojlim_n K_2(H_n)^{\wedge} & \xrightarrow{cl} & \varprojlim_n H^2(H_n, \mathbb{Z}/p^n \mathbb{Z}(2)) & \xrightarrow{C_1} & H^2(H_1, \mathbb{Z}_p(1)) \\ [p^{n-1}]_* \downarrow & & \text{Cor}_{H_n/H_1}(\zeta_n) \downarrow & & \text{id} \downarrow \\ \varprojlim_n K_2(H_1(\zeta_n))^{\wedge} & \xrightarrow{cl} & \varprojlim_n H^2(H_1(\zeta_n), \mathbb{Z}/p^n \mathbb{Z}(2)) & \xrightarrow{C_2} & H^2(H_1, \mathbb{Z}_p(1)) \end{array}$$

Here $C_1 = \text{Cor}_{H_n/H_1} \circ \cup(\zeta_n)^{-1}$ and $C_2 = \text{Cor}_{H_1(\zeta_n)/H_1} \circ \cup(\zeta_n)^{-1}$. By this diagram it follows that if $\partial((\beta'_n)_n) \neq 0$, then $cl([p^{n-1}]_* \beta'_n) \neq 0$ for sufficiently large n .

It implies that $0 \neq cl([\pi^{n-1}]_* \alpha'_n) \in H^2(F_n \mathfrak{P}(E), \mathbb{Z}/p^n \mathbb{Z}(2))$ where $F_n \mathfrak{P}$ is completion of F_n by the prime over $\mathfrak{P}$.

Consider the restriction of $\tilde{cl}([\pi^{n-1}]_* \alpha'_n)$ to $H^1(F_n \mathfrak{P}, T_p(E)(1))$. We need a splitting $H^1(F_n \mathfrak{P}, T_p(E)(1)) \rightarrow H^2(E_{F_n \mathfrak{P}}, \mathbb{Z}/p^n \mathbb{Z}(2))$ and the following lemma enables us to define such a splitting.

Lemma 3.2. $\sharp H^2(F_n \mathfrak{P}, \mathbb{Z}/p^n \mathbb{Z}(2)) = \sharp H^0(F_n \mathfrak{P}, \mathbb{Z}/p^n \mathbb{Z}(-1))$ is bounded with respect to n .

Proof. Considering the Weil pairing $E[\pi^n] \times E[\bar{\pi}^n] \rightarrow \mu_{p^n}$, we see that if $F_n \mathfrak{P} \supset F_{\mathfrak{P}}(\mu_{p^m})$ for a number $m \leq n$, then $F_n \mathfrak{P} \supset F_{\mathfrak{P}}(E[\bar{\pi}^m])$. Since $\mathfrak{P}$ is finitely decomposed in $F(E[\bar{\pi}^{\infty}])$, the residue field of $F_{\mathfrak{P}}(E[\bar{\pi}^{\infty}])$ is $\overline{\mathbb{F}_p}$. As $F_n \mathfrak{P}$ is totally ramified over $F_{\mathfrak{P}}$, it follows that m is bounded. $\square$

By this lemma we have a number M which is a power of p and a map

$$u_n : H^1(F_n \mathfrak{P}, T_p(E)(1)) \rightarrow H^2(E_{F_n \mathfrak{P}}, \mathbb{Z}/p^n \mathbb{Z}(2))$$

for each $n \geq 1$ such that

$$u_n(\tilde{cl}([\pi^{n-1}]_* \alpha'_n) = Mcl([\pi^{n-1}]_* \alpha'_n).$$

Noting that the action of the Frobenius Φ on $E[a]$ is by π , one can show the following in a similar way to the proof of Theorem 2.4.

Theorem 3.3. *For $\gamma \in E[a]$, set $g_a^\gamma := T_\gamma^* g_a$. Then we have the equality*

$$h^{-1}((\beta'_n)) = B' := [F'_0 : F][p]_* \sum_{\gamma \in E[a]} \{\theta^* g_a^\gamma, \theta^* g_a^\gamma(s[+]_{\widehat{\mathbb{G}}_{mW}} \epsilon - 1)\}.$$

Since $d \log$ and the pull back $[p]^*$ commutes on S' , it also commutes on $K_2(S')$. As

$$\begin{aligned} [p]^* B' &= [F'_0 : F] \sum_{\zeta^p=1} \sum_{\gamma \in E[a]} \\ &\quad \{\theta^* g_a^\gamma(s[+]_{\widehat{\mathbb{G}}_{mW}} \zeta - 1), \theta^* g_a^\gamma(s[+]_{\widehat{\mathbb{G}}_{mW}} \zeta - 1[+]_{\widehat{\mathbb{G}}_{mW}} \epsilon - 1)\} \end{aligned}$$

we see that

$$\begin{aligned} &[p]^* \partial((\beta'_n)) \\ &= \frac{[F'_0 : F]}{p} \Phi^{-1} \left(\sum_{\zeta^p=1} \sum_{\gamma \in E[a]} D \log \theta^* g_a^\gamma(s[+]_{\widehat{\mathbb{G}}_{mW}} \zeta - 1) \right. \\ &\quad \left. \times D \log \theta^* g_a^\gamma(s[+]_{\widehat{\mathbb{G}}_{mW}} \zeta - 1[+]_{\widehat{\mathbb{G}}_{mW}} \zeta_1 - 1) \right) d \log q. \end{aligned}$$

We write the last term as $\frac{[F'_0 : F]}{p} \Phi^{-1} G(s) d \log q$.

Let $\lambda : \widehat{E} \rightarrow \widehat{\mathbb{G}}_a$ be a logarithm defined over F and let z be the parameter of $\widehat{\mathbb{G}}_a$ such that $z = \lambda(t) = t + \dots$. We have the equality

$$\Omega_p \frac{d}{dz} \circ \lambda_* \theta_* = \lambda_* \theta_* \circ D$$

where $\Omega_p \in W^*$ is such that $t = \theta(s) = \Omega_p s + \dots$. Let $\xi : E(\mathbb{C}) \rightarrow \mathbb{C}/L$ be the inverse of the analytic uniformization associated to dz . Here L is the period lattice of dz . For a point $v \in E[\pi]$, $\frac{d}{dz} \lambda_* g_a^\gamma(z[+]_{\widehat{\mathbb{G}}_a} \lambda(v))$ is the Taylor expansion of $\frac{d}{dz} \xi_* g_a^\gamma(z + \xi(v))$ at $z = 0$. Hence by looking at $G(s)$ we obtain the following corollary.

Corollary 3.4.

If the function

$$\sum_{v \in E[\pi]} \sum_{\gamma \in E[a]} \frac{d}{dz} \log(\xi_* g_a^\gamma)(z + \xi(v)) \times \frac{d}{dz} \log(\xi_* g_a^\gamma)(z + \xi(v + v_1))$$

on $\mathbb{C}/L$ is nonzero, then $\tilde{cl}([\pi^{n-1}]_ \alpha'_n)$ is nonzero for sufficiently large n .*

Appendix: Euler Systems. Let $\mathcal{S}$ be the set of ideals of O_K which are divisible only by primes which split completely in F and which are relatively prime to $\mathfrak{f}_a$. For each $\mathfrak{m} \in \mathcal{S}$, Let $F_{\mathfrak{m}} := F(E[\mathfrak{m}])$. Fix a system of torsion points $(v_{\mathfrak{m}})_{\mathfrak{m} \in \mathcal{S}}$ such that

(1) $v_{\mathfrak{m}}$ is a generator of $E[\mathfrak{m}]$ for each $\mathfrak{m} \in \mathcal{S}$.

(2) If $\mathfrak{m}_2 = \mathfrak{m}_1 \mathfrak{n}$ with $\mathfrak{m}_2 \in \mathcal{S}$, then $v_{\mathfrak{m}_1} = \phi(\mathfrak{n})v_{\mathfrak{m}_2}$.

We set $\alpha_{\mathfrak{m}} := \{g_a, T_{v_{\mathfrak{m}}}^* g_a\} \in K_2(F_{\mathfrak{m}}(E))$ and set $\alpha'_{\mathfrak{m}} := [\phi(\mathfrak{m})]_* \sum_{\gamma \in E[a]} T_{\gamma}^* \alpha_{\mathfrak{m}}$. Then it

is seen that $\alpha'_{\mathfrak{m}} \in \Gamma(E_{F_{\mathfrak{m}}}, K_2)$ in the similar way to the argument in the beginning of §3. So that there is a class $\tilde{cl}(\alpha'_{\mathfrak{m}}) \in H^1(F_{\mathfrak{m}}, T_p(E)(1))$. In the similar way to the proof of (EN1) in ([Sch], Appendix) one obtains the following result:

Proposition A.1. *Let $\mathfrak{m} \in \mathcal{S}$ and assume that $\mathfrak{l} \nmid \mathfrak{m}$ for a prime $\mathfrak{l}$. Then the following equality holds:*

$$N_{F_{\mathfrak{m}}/F_{\mathfrak{m}/\mathfrak{l}}} \alpha'_{\mathfrak{m}} = \begin{cases} \alpha'_{\mathfrak{m}/\mathfrak{l}} & \mathfrak{l}^2 \mid \mathfrak{m} \\ (1 - [\phi(\mathfrak{l})]_* Fr_{\mathfrak{l}}^{-1}) \alpha'_{\mathfrak{m}/\mathfrak{l}} & \mathfrak{l}^2 \nmid \mathfrak{m}. \end{cases}$$

The classes $\tilde{cl}(\alpha'_{\mathfrak{m}})$ satisfy the same relation as $\alpha'_{\mathfrak{m}}$. This implies the classes $\tilde{cl}(\alpha'_{\mathfrak{m}})$ form an Euler system.

REFERENCES

- [dS] de Shalit, E., *Iwasawa theory of elliptic curves with complex multiplication*, Academic press, 1987.
- [Fu] Fukaya, T., *The theory of Coleman power series for K_2* , to appear in the Journal of Alg.Geom. (2000).
- [Ga] Garland, H., *A finiteness theorem for K_2 of a number field*, Advances in Math. **94**(2) (1971), 534–538.
- [Gr] Grothendieck, A., *EGA IV (Premiere Partie)*, Publ. Math. I.H.E.S. **20** (1964).
- [Hy] Hyodo, O., *On the Hodge-Tate decomposition in the imperfect residue field case*, Journal für die Reine und Ang. Math. **365** (1986), 97–113.
- [Qu] Quillen, D., *Higher algebraic K-theory I*, Springer Lecture notes in Math. **341** (1973), 85–147.
- [Ru] Rubin, K., *The “main conjecture” of Iwasawa theory for imaginary quadratic fields*, Invent. Math. **103** (1991), 25–68.
- [Sch] Scholl, A.J., *An introduction to Kato’s Euler systems*, London Math. Soc. Lect. Notes **254** (1998), pp. 379–460.